

Conferences and Reviews

Blood-Brain Barrier and New Approaches to Brain Drug Delivery

Moderator

WILLIAM M. PARDRIDGE, MD

Discussants

RUBEN J. BOADO, PhD; KEITH L. BLACK, MD; and PASQUALE A. CANCELLA, MD

This is an edited summary of an Interdepartmental Conference arranged by the Department of Medicine of the UCLA School of Medicine, Los Angeles. The Director of Conferences is Anthony T. Campagnoni, PhD, Professor of Psychiatry and Biobehavioral Sciences.

Morbidity caused by brain dysfunction affects more than 50 million persons in the United States. Although new neuropharmaceuticals have the potential for treating specific brain diseases, they may not effectively enter brain from blood. Safe strategies are needed for drug delivery through the brain capillary wall, which makes up the blood-brain barrier in vivo. Two of these strategies are reviewed, as are related new developments in the molecular and cell biology of the brain capillary endothelium. The production of chimeric peptides represents a physiologic-based strategy for drug delivery. It entails the covalent coupling of the neuropharmaceutical to a brain transport vector, allowing transportation through the blood-brain barrier. Another strategy is biochemical opening of the blood-brain barrier: intracarotid leukotriene infusion is a method for selectively increasing blood-brain barrier permeability in brain tumors without affecting barrier permeability in normal brain tissue.

(Partridge WM, Boado RJ, Black KL, Cancelli PA: Blood-brain barrier and new approaches to brain drug delivery. *West J Med* 1992 Mar; 156:281-286)

WILLIAM M. PARDRIDGE, MD*: The morbidity of cancer and heart disease is dwarfed by illness caused by disorders of the brain, which affect 50 to 100 million persons in the United States (Table 1). The number of disorders of the brain commonly seen in the practice of internal medicine may be expanded even further because such problems as peptic ulcer, hypertension, or irritable bowel syndrome may be amenable to treatment by direct drug action on the brain.¹ In the 1990s, the Decade of the Brain, the traditional trial and error approach to drug development and design is giving way to rational drug design based on three-dimensional models of receptor structure. For example, new drugs for treating depression, stroke, Alzheimer's disease, fever, or anxiety may include corticotrophin-releasing hormone antagonists, calcitonin gene-related peptide agonists, amyloid protease inhibitors, α -melanocyte-stimulating hormone agonists, or cholecystokinin receptor antagonists, respectively.¹ In future, the driving force of drug design may be the expanding base of neuropeptide receptor biology that has emerged from research.

Unfortunately, drugs developed from rational drug design based on considerations of receptor biology may not be effective neuropharmaceuticals if these drugs are not transportable across the brain capillary wall, which makes up the blood-brain barrier in vivo. For example, the first product of recombinant DNA technology that may be used for treating a neurologic disease is nerve growth factor for treating Alzhei-

mer's disease.² Nerve growth factor is a large protein of approximately 25,000 molecular weight and is not known to cross the blood-brain barrier.³ Although proposals have been made to administer nerve growth factor through intraventricular drug delivery,⁴ this approach requires a neurosurgical procedure. Moreover, the two caveats of intraventricular drug delivery are: the drug is distributed only to the surface of the brain and not deep into brain parenchyma,⁵ and the drug is readily distributed into the peripheral circulation owing to rapid efflux of drug from the ventricular cerebrospinal

TABLE 1.—Common Brain Disorders in the United States*

Disorder	No. of Affected Persons, in millions
Migraine headache	25
Alcohol abuse	25
Anxiety or phobia	25
Sleep disorders	20
Depression or mania	12
Drug abuse	10
Obsessive-compulsive disorder	4
Alzheimer's disease	3
Schizophrenia	2
Stroke	2
Epilepsy	2
Human immunodeficiency virus infection	1.5
Parkinson's disease	.5

*From Partridge.¹

*Professor, Department of Medicine, UCLA School of Medicine.

This study was supported in part by National Institutes of Health Program Project NS-25554.

Reprint requests to William M. Partridge, MD, Department of Medicine, UCLA School of Medicine, Los Angeles, CA 90024.

ABBREVIATIONS USED IN TEXT

cDNA = complementary DNA
mRNA = messenger RNA
pⁱ = isoelectric point

fluid compartment to the bloodstream through the arachnoid granulations at the superior sagittal sinus.⁶ Another traditional approach for drug delivery are liposomes.⁷ Liposomes, however, are too large to traverse the blood-brain barrier and have not been proved to be effective neuropharmaceutical delivery vehicles.¹

The importance of developing effective neuropharmaceuticals and the lack of efficacy of existing strategies for drug delivery to brain emphasize the need to develop new approaches to drug transport across the blood-brain barrier. Two approaches discussed in this review have emanated as practical applications of fundamental basic blood-brain barrier research. The first approach is using chimeric peptides as a vehicle for neuropharmaceutical delivery through the blood-brain barrier. The second is biochemical opening of the blood-brain barrier.

Chimeric Peptides

Chimeric peptides are formed when a nontransportable pharmaceutical peptide—for example, β -endorphin for the treatment of pain—is coupled to a blood-brain barrier transport vector.⁸ The latter is a peptide or plasma protein that normally enjoys receptor-mediated or absorptive-mediated transcytosis through the blood-brain barrier. Examples of compounds that traverse the blood-brain barrier by receptor-mediated transcytosis include insulin,⁹ transferrin,¹⁰ a monoclonal antibody to surface epitopes of the transferrin receptor,¹¹ and possibly insulin-like growth factors.¹² Proteins that traverse the blood-brain barrier via absorptive-mediated transcytosis include polycationic proteins, such as histone,¹³ cationized albumin,¹⁴ cationized antibodies,¹⁵ or lectins, such as wheat germ agglutinin.¹⁶ Although native albumin, which is an acidic protein with an isoelectric point (pⁱ) of 4 to 5, does not normally cross the blood-brain barrier, chemical modification of the protein, which results in the formation of cationized albumin, pⁱ = 8 to 9, causes the protein to undergo absorptive-mediated transcytosis through the blood-brain barrier.^{14,17} More recent studies have shown that an antitransferrin receptor antibody¹¹ undergoes receptor-mediated transcytosis through the blood-brain barrier at rates that are severalfold faster than the transport of cationized albumin.¹⁸

Proteins such as cationized albumin or antitransferrin receptor monoclonal antibodies may be used as a vector to ferry across the blood-brain barrier compounds that are pharmacologically active in brain (Figure 1).¹⁹ The four steps of the chimeric peptide transport process are receptor-mediated endocytosis at the blood side of the blood-brain barrier; receptor-mediated exocytosis of the chimeric peptide at the brain side of the blood-brain barrier with release into brain interstitial fluid; thiol-based cleavage of the disulfide bond adjoining the chimeric peptide, which results in the release within brain of pharmacologically active peptide; and binding of the pharmacologically active peptide to its cognate receptor in brain after release from the vector. Previous studies have provided evidence that a β -endorphin cationized albumin chimeric peptide undergoes absorptive-mediated endocytosis




Figure 1.—The delivery of chimeric peptide through the blood-brain barrier is viewed as a process composed of 4 overall steps (see text). A is a transportable vector, for example, antitransferrin receptor monoclonal antibody; B, non-transportable (pharmacologically active) peptide, for example, β -endorphin; A-R, peptide A, for example, transferrin receptor; B-R, peptide B receptor (from Pardridge et al¹⁹; reproduced with permission from *Endocrinology*).

and exocytosis at the blood-brain barrier in vivo. Also, the brain contains the necessary disulfide reductases that are sufficiently active to cleave the chimeric peptide, which results in the release of free β -endorphin from its cationized albumin vector.¹⁹ Cationized albumin has proved to be an effective blood-brain barrier transport vector for experimental studies in laboratory animals. The use of this vector in humans may be limited by the rapid renal uptake of cationized albumin, however.¹⁷ While highly cationic proteins are known to be nephrotoxic,²⁰ recent studies have shown that mildly cationized homologous albumins may be administered repetitively to laboratory animals without measurable nephrotoxicity.¹⁷ Nevertheless, it would be advantageous to develop vectors with a higher degree of brain specificity. Developing such vectors may invariably arise from a molecular biologic analysis of brain capillary-specific proteins. The production of monoclonal antibodies to brain capillary-specific proteins may prove to be brain transport vectors with a high degree of brain specificity.

Blood-Brain Barrier Glucose Transporter, a Model Brain Capillary-Enriched Protein

RUBEN J. BOADO, PhD*: A vector with a relatively high degree of brain specificity is a monoclonal antibody to the transferrin receptor.¹¹ The transferrin receptor is a brain capillary-enriched protein,²¹ that is, a protein found in high concentration in brain capillary endothelium relative to neurons or glial cells or to microvascular endothelium in peripheral tissues. Other brain capillary-enriched proteins are listed in Table 2.^{1,21-33} When sections of animal or human brain are stained with antibodies to these proteins, the microvascular endothelium is markedly illuminated owing to the high concentration of these proteins in brain capillary endothelium. The brain capillary-enriched proteins, however, may also be localized in peripheral tissues. In contrast, brain capillary-specific proteins are thought to be expressed only in the brain capillary endothelium and not in peripheral tissues, and two brain capillary-specific proteins have been identified (Table 2).

The identification and molecular cloning of brain capillary-specific protein genes may lead to the development of monoclonal antibodies specific to these proteins. The molecular cloning of brain capillary-specific protein genes requires the development of brain capillary complementary DNA

*Assistant Professor, Department of Medicine, UCLA School of Medicine.

TABLE 2.—Brain Capillary-Enriched Proteins and Brain Capillary-Specific Proteins*

Functional Class	Protein
Brain Capillary-Enriched Proteins	
Nutrient transporters.....	GLUT-1 glucose transporter Low Km NAA transporter
Peptide transporters.....	Transferrin receptor Insulin receptor
Drug transporters.....	P-glycoprotein (MDR gene product)
Vitamin transporters.....	Cellular retinol binding protein
Cholesterol transporters.....	Apolipoprotein A-I
Enzymes.....	γ -Glutamyltranspeptidase Aminopeptidase M
Phosphoproteins.....	50-55 KDa triplet
HLA antigens.....	Class I (endothelium) DR-antigen (pericyte, smooth muscle)
Surface antigens.....	CD36 HT7 140 KDa pericyte protein 25 KDa triplet
Brain Capillary-Specific Proteins	
Surface antigens.....	45 and 53 KDa proteins
MDR = multidrug resistance, NAA = neutral amino acid	
*From Pardridge, ¹ compiled from data in references 21-33.	

(cDNA) libraries. It is probably not possible to isolate cDNA clones corresponding to brain capillary-specific protein genes by screening cDNA libraries made from total brain. The volume of the brain capillary endothelial cell is only about 1 μ l per gram of brain. There is only approximately 1 ml of endothelial cytoplasm in an entire human brain and only 1 μ l of endothelial cytoplasm in an entire rat brain.¹ Therefore, less than 0.2% of the cDNA clones in a total brain library may be derived from reverse transcription of brain capillary-derived messenger RNA (mRNA). Consequently, it is necessary to isolate mRNA density from brain capillaries, which may be isolated from either laboratory animal or human autopsy brain (Figure 2).³⁴ Messenger RNA has been directly isolated from either bovine or pig brain capillaries, and blood-brain barrier cDNA libraries have been prepared in λ gt10³² or λ gt11 vectors.³³ From these libraries, several brain capillary-enriched protein gene cDNAs have been cloned, including those for γ -glutamyl transpeptidase,²⁴ the GLUT-1 glucose transporter,^{33,35} and apolipoprotein A-I.³²

The blood-brain barrier GLUT-1 glucose transporter may be a brain capillary-enriched protein target for drug delivery because recent studies have shown the concentration of this protein in brain microvessels is particularly high, for example, 10 picomol per mg of protein.³⁶ Thus, the concentration of the GLUT-1 glucose transporter is approximately 100-fold more abundant than is the transferrin receptor.³⁷ In addition, the GLUT-1 glucose transporter protein is expressed selectively in brain microvascular endothelium with minimal, if any, expression in neurons or glial cells. Recent quantitative Western blotting, in situ hybridization, quantitative Northern blotting, and quantitative electron microscopic immunogold studies have all indicated that the GLUT-1 glucose transporter is selectively expressed in brain microvascular endothelium with no measurable expression in neurons or glial cells.^{36,38,39} The GLUT-1 glucose transporter protein, however, is also expressed in peripheral tissues, including perivascular hepatocytes, renal collecting duct epithelium, peripheral nerve perineurium, syncytiotrophoblast of the placenta, placenta-fetal microvascular endothelium, and tes-




Figure 2.—A light micrograph shows capillaries isolated from a human brain at autopsy. The microvessels are comprised of endothelial cells and pericytes (in an approximate 3:1 ratio) encased in a common basement membrane (magnification $\times 500$; from Pardridge et al³⁴; reproduced with permission from *Journal of Neurochemistry*).

ticular microvascular endothelium.⁴⁰ Therefore, although the GLUT-1 transporter protein is specific to endothelium with respect to neurons or glial cells, it is also found in various epithelial and endothelial cells in peripheral tissues.

Biochemical Opening of the Blood-Brain Barrier—Leukotrienes

KEITH L. BLACK, MD*: Capillaries within primary tumors contain a blood-tumor barrier that impairs delivery of antitumor compounds to neoplastic cells within the brain.⁴¹⁻⁴³ The inability to deliver antineoplastic drugs across the blood-tumor barrier has led to experimental attempts to open the barrier.^{44,45} One approach has used intracarotid infusion of mannitol, which causes a transient osmotic shrinkage of the brain capillary endothelial cells and disrupts the blood-brain barrier tight junctions in normal brain, as well as in brain tumors.^{43,45-47} The osmotic barrier disruption approach nonspecifically increases the delivery of chemotherapeutic compounds and monoclonal antibodies to brain.⁴⁴ Osmotic disruption of the blood-brain barrier, however, results in only a modest increase in drug levels within the actual tumor, compared with the relatively large increase in drug levels in normal brain.⁴⁸ Therefore, this technique increases the exposure of the normal brain parenchyma to the deleterious effects of the chemotherapeutic compounds,⁴⁶ whereas suboptimal drug levels may actually reach the tumor. We recently described an experimental method that circumvents this problem and uses leukotrienes to selectively open the barrier within the tumor while leaving the normal blood-brain barrier intact.⁴⁹

The rate-limiting step in leukotriene production is the 5-lipoxygenase enzyme. Brain lipids are rich in membrane-bound arachidonic acid. During a variety of metabolic events, arachidonic acid is released from cell membranes by phospholipase A2 or phospholipase C. Free arachidonic acid, once released, is either reincorporated into the cell membrane or oxidized by one of two pathways: the cyclooxygenase pathway to prostaglandins or the 5-lipoxygenase pathway to leukotrienes. The 5-lipoxygenase pathway—requiring molecular oxygen and calcium—generates the hydroxyleuko-

*Assistant Professor, Department of Surgery, Division of Neurosurgery, UCLA School of Medicine.

triene B4 and the peptidoleukotrienes C4, D4, E4, and F4 via the epoxide intermediate leukotriene A4. The blood-brain barrier γ -glutamyl transpeptidase (Table 2) inactivates leukotriene C4 to leukotriene D4 and leukotriene E4 to leukotriene F4.⁵⁰ This enzyme is unique to brain capillaries as it is not present in systemic capillaries.⁵¹

Leukotrienes are quite potent in their ability to increase vascular permeability in systemic (noncerebral) capillaries.⁵² Normal brain capillaries, however, resist the efforts of leukotrienes to increase their vascular permeability. In effect, normal brain capillaries seem to contain an "enzymatic barrier" that inactivates vasoactive compounds such as leukotriene C4. Injured brain capillaries or capillaries in brain tumors lose their "enzymatic barrier" and show increased vascular permeability after leukotriene C4 infusions.⁴⁹

In human brain tumors, a correlation between brain edema and tissue levels of leukotriene C4 has been shown.⁵³ An index of peritumoral edema was estimated from preoperative contrast-enhanced computed tomographic scans. There was a significant correlation between edema and the log leukotriene C4 levels, with $P < .003$ in all patients. Metastatic tumors had the highest leukotriene C4 level at 13.8 ± 8.5 pg per mg of tissue (mean \pm standard error) and the highest index of edema at 5.7 ± 1.8 . The mean leukotriene C4 level in gliomas was 6.2 ± 2.3 pg per mg of tissue and the edema index was 2.1 ± 0.6 . There was no edema, no tumor in the temporal lobes removed for seizure, and their level of leukotriene C4 was 0.4 ± 0.1 pg per mg of tissue. These data show that the formation of leukotriene C4 is stimulated by intracranial tumors.

In experimental brain tumors in rats (C6 glioma, RG-2 glioma, and Walker 256 metastatic tumor) γ -glutamyl transpeptidase could not be shown within tumor capillaries.⁴⁹ Because brain tumors stimulate leukotriene C4 formation, and tumor capillaries—unlike normal brain capillaries—lack γ -glutamyl transpeptidase to inactivate leukotriene C4, leukotrienes could play an important role in the development of vasogenic edema surrounding brain tumors and in increasing capillary permeability within tumors.

It is important for delivery of antitumor compounds to have the ability to further open the capillary barrier within tumors. Recently intracarotid infusions of leukotriene C4 were shown to selectively open the blood-tumor barrier in rats with RG-2 gliomas.⁴⁹ Blood-brain and blood-tumor permeability was determined by quantitative autoradiography using ^{14}C aminoisobutyric acid. Leukotriene C4 (total dose, 4 μg) infused into the carotid artery ipsilateral to the tumor increased twofold the unidirectional transfer constant for permeability within the tumor, whereas no effect on permeability was seen in normal brain. No γ -glutamyl transpeptidase activity was seen in tumor capillaries, in contrast to high γ -glutamyl transpeptidase in normal brain capillaries. These findings support the hypothesis that normal brain capillaries resist the vasogenic effects of leukotriene C4, whereas leukotriene C4 may increase permeability in tumor capillaries.

Loss of γ -glutamyl transpeptidase is not, however, limited in brain to tumor capillaries. γ -Glutamyl transpeptidase activity is also decreased in capillaries in ischemic tissue at 48 and 72 hours after middle cerebral artery occlusion in rats.⁵⁴ In comparison, high γ -glutamyl transpeptidase in normal brain capillaries and moderate γ -glutamyl transpeptidase in capillaries in ischemic tissue 24 hours after middle cerebral artery occlusion was reported. At 72 hours after

middle cerebral ischemia, the infusion of leukotriene C4 (total dose, 4 μg) into the carotid artery ipsilateral to the ischemia selectively increased blood-brain barrier permeability approximately threefold within core ischemic tissue and tissue adjacent to the ischemic core.⁵⁴ No effect on blood-brain barrier permeability was seen within nonischemic brain tissue or in ischemic tissue after only 24 hours of ischemia.

Acivicin is an inhibitor of γ -glutamyl transpeptidase.⁵⁴ When rats are pretreated with acivicin, the ability of leukotriene C4 to increase blood-brain barrier permeability in ischemic brain even at 24 hours is significantly enhanced. At 24 hours, when only a moderate reduction in γ -glutamyl transpeptidase has occurred, further blocking of γ -glutamyl transpeptidase activity with acivicin results in a significant enhancement of the ability of leukotriene C4 to open the blood-brain barrier.⁵⁴

These experiments show that brain capillaries contain an "enzymatic barrier" that inactivates vasoactive compounds, like leukotrienes, which increase blood-brain barrier permeability. In some pathologic conditions—that is, brain tumors, ischemia—this "enzymatic barrier" is lost. One component of the enzymatic barrier seems to be γ -glutamyl transpeptidase. Its loss in tumor capillaries and in ischemia permits leukotriene C4 to increase vascular permeability in these conditions and produce vasogenic edema. Because leukotriene C4 production is significantly increased in brain tumors, leukotrienes may play an important role in edema surrounding brain tumors. Clinically important, the selective vulnerability of tumor capillaries to leukotrienes can be used therapeutically. Because tumor capillaries in brain tumors form a barrier to many chemotherapeutic compounds and monoclonal antibodies, leukotriene C4 may be used to open selectively the barrier within tumors while leaving the normal blood-brain barrier intact. One important advantage of selective barrier opening would be protecting normal brain tissue from the potentially deleterious effects of antineoplastic compounds.

Biochemical opening of the blood-brain barrier by vasoactive substances such as leukotrienes may involve cell-to-cell interactions at the cerebral microcirculation. The latter is comprised of at least four cell types: endothelium, which constitutes the permeability barrier; astrocyte foot processes, which invest more than 95% of the surface of the brain capillary; pericytes, which have contractile and phagocytic functions at the microcirculation; and nerve endings, which directly innervate brain microvascular endothelium.¹ To understand the mechanisms of cell-to-cell interactions at the brain microcirculation, in vitro tissue culture models have been established to investigate the cell biology of endothelium-astrocyte interactions.

Cell Biology of Brain Endothelium and Astrocyte Interactions

PASQUALE A. CANCELLA, MD*: Our laboratory has used in vivo and in vitro systems to explore the complex interactions that occur between endothelial cells and astrocytes and that lead to the development of properties associated with the blood-brain barrier. The in vivo system most often used by us has been the localized application of liquid nitrogen to the intact skull of the mouse to produce a discrete freeze injury.⁵⁵

*Professor, Department of Pathology, UCLA School of Medicine.

This area of injury undergoes a reproducible series of changes including necrosis, monocyte-macrophage infiltration, endothelial proliferation and neovascularization, and astrocyte proliferation and migration that lead to restitution of endothelial-astrocyte relations and restoration of the blood-brain barrier. Our studies used ^3H -thymidine as a marker for the initiation of DNA synthesis in endothelial cells and astrocytes at intervals after the injury.^{56,57} Both endothelial cells and astrocytes are normally quiescent cells that quickly return to the mitotic pool in response to injury. This coincides with the migration of monocyte-macrophage into the area of injury from the circulation. Interruption of the monocyte-macrophage response reduces the endothelial response to approximately 10%. Thus, factors derived from the monocyte-macrophage are operational in the endothelial response. Cytokines may also be involved in restoring the endothelium of the injured vessels and in neovascularization because in-vitro studies have shown a chemotactic effect of monocyte-macrophage on endothelium.⁵⁸

The astrocytes adjacent to the area of injury undergo a similar ^3H -thymidine incorporation after an injury, but the maximum response is of lower magnitude (10%) and occurs later—day 5 compared with day 3 to 4 after injury—than the endothelial cell response. In vitro studies have shown that cytokines derived from monocyte-macrophage may be operational in this response and that endothelial cells may liberate chemotactic factors, which attract astrocytes from the uninjured brain adjacent to the injured area and to the new vessels.⁵⁸ Studies using the freeze-injury model have shown that endothelial cells in vessels adjacent to the injury initially have the enzyme γ -glutamyl transpeptidase, but within five to seven days of the injury, the enzyme is no longer found in the endothelium.⁵⁷ This corresponds to the time when the endothelial cells migrate into the area of injury as part of the process of neovascularization and correlates with the loss of contact of the astrocyte with the endothelium.⁵⁶ When the astrocyte reestablishes contact with the endothelial cell, γ -glutamyl transpeptidase is found once again in the endothelium.⁵⁷ This supports the in vivo findings that interaction of the astrocyte with the endothelial cell is necessary for the induction and maintenance of γ -glutamyl transpeptidase expression and that a factor liberated by the astrocyte is responsible for this effect.^{59,60}

Endothelial cells influence responses in the astrocyte. Media conditioned by cerebral endothelial cells contain a factor or factors that stimulate the incorporation of ^3H -thymidine by the astrocyte.⁶¹ This factor is a peptide of greater than 50,000 molecular weight that is destroyed by protease digestion and is different from other known growth factors. Rather than stimulating ^3H -thymidine incorporation in endothelial cells through an autocrine effect, the factor does not affect endothelial cells.

Dr Takashi Minakawa, in our laboratory, has used his model of developing vascular-like structures in vitro into a model replicating the relation of the vascular endothelium and astrocytes.⁶² The system used gelatin-coated slides in chambers as a base for developing branching tube-like structures when coated with vitrogen. Astrocytes or pericytes added to the surface of the vitrogen migrate through this layer and develop contacts with the endothelial structures. Astrocytes and pericytes respond differently. By 24 hours after the random addition of the pericytes to the system, 100% of the cells are in contact with the endothelium. Astrocytes are

slower to respond because 10% are in contact by 5 hours, 75% are associated with the vessels at 24 hours, and 93% by 6 days.

A chemotactic assay has been developed using several features from this system.⁶² Use of it has shown that these vascular-like structures, but not monolayer cultures of endothelium, release a chemotactic factor involved in the attraction of the astrocyte to the endothelium. This factor is liberated from the better-differentiated cells forming the vascular tubes and suggests that the response is coordinated and may serve to attract the astrocyte to the endothelium, as was suggested by the in vivo model.

The studies described in this presentation represent some, but not all, of the types of studies that have been done to show that the cell-to-cell interaction between endothelium and astrocyte is important in establishing and maintaining the blood-brain barrier.⁶³⁻⁶⁸ Many investigators are interested in this area, so we may soon see important and detailed reports describing not only the cell biology but also the molecular mechanisms of these complex and fascinating relations.

Ongoing basic research into the molecular and cell biology of the blood-brain barrier is expanding further the definition of brain capillary-specific proteins and the molecular mechanisms involved in paracrine cellular interactions at the level of the cerebral microcirculation. This research invariably provides insights into the pathogenesis of brain diseases.⁶³ Basic research in blood-brain barrier molecular and cell biology has also led to new approaches to brain drug delivery. As molecular and neurobiologic research further defines the mechanisms of disorders of the brain, new therapies may emerge. An essential component for developing neuropharmaceuticals of the future will be integrating these with the emerging paradigms of blood-brain barrier drug delivery.

REFERENCES

1. Pardridge WM: Peptide Drug Delivery to the Brain. New York, Raven Press, 1991
2. Marx J: NGF and Alzheimer's: Hopes and fears. *Science* 1990; 247:408-410
3. Levi-Montalcini R, Angeletti PU: Nerve growth factor. *Physiol Rev* 1968; 48:534-569
4. Harbaugh RE: Nerve growth factor as a potential treatment in Alzheimer's disease. *Biomed Pharmacother* 1989; 43:483-485
5. Blasberg RG, Patlak C, Fenstermacher JD: Intrathecal chemotherapy: Brain tissue profiles after ventriculocisternal perfusion. *J Pharm Exp Ther* 1975; 195:73-83
6. Aird RB: A study of intrathecal, cerebrospinal fluid-to-brain exchange. *Exp Neurol* 1984; 86:342-358
7. Gregoriadis G: The carrier potential of liposomes in biology and medicine. *N Engl J Med* 1976; 295:704-710
8. Pardridge WM: Receptor-mediated peptide transport through the blood-brain barrier. *Endocr Rev* 1986; 7:314-330
9. Duffy KR, Pardridge WM: Blood-brain barrier transcytosis of insulin in developing rabbits. *Brain Res* 1987; 420:32-38
10. Fishman JB, Rubin JB, Handrahan JV, Connor JR, Fine RE: Receptor-mediated transcytosis of transferrin across the blood-brain barrier. *J Neurosci* 1987; 18:299-304
11. Friden PM, Walus L, Musso GF, Taylor MA, Malfroy B, Starzyk RM: Anti-transferrin receptor antibody and antibody-drug conjugates cross the blood-brain barrier. *Proc Natl Acad Sci USA* 1991; 88:4771-4775
12. Duffy KR, Pardridge WM, Rosenfeld RG: Human blood-brain barrier insulin-like growth factor receptor. *Metabolism* 1988; 37:136-140
13. Pardridge WM, Triguero D, Buciak JL: Transport of histone through the blood-brain barrier. *J Pharmacol Exp Ther* 1989; 251:821-826
14. Kumagai AK, Eisenberg J, Pardridge WM: Absorptive-mediated endocytosis of cationized albumin and a β -endorphin-cationized albumin chimeric peptide by isolated brain capillaries. Model system of blood-brain barrier transport. *J Biol Chem* 1987; 262:15214-15219
15. Triguero D, Buciak JL, Pardridge WM: Blood-brain barrier transport of cationized immunoglobulin G. Enhanced delivery compared to native protein. *Proc Natl Acad Sci USA* 1989; 86:4761-4765
16. Broadwell RD, Balin BJ, Salzman M: Transcytotic pathway for blood-borne protein through the blood-brain barrier. *Proc Natl Acad Sci USA* 1988; 85:632-636
17. Pardridge WM, Triguero D, Buciak JL, Yang J: Evaluation of cationized rat

albumin as a potential blood-brain barrier drug transport vector. *J Pharmacol Exp Ther* 1990; 255:893-899

18. Pardridge WM, Buciak JL, Friden P: Selective transport of anti-transferrin receptor antibody through the blood-brain barrier in vivo. *J Pharmacol Exp Ther* 1991; 259:66-70

19. Pardridge WM, Triguero D, Buciak JL: β -Endorphin chimeric peptides: Transport through the blood-brain barrier in vivo and cleavage of disulfide linkage by brain. *Endocrinology* 1990; 126:977-984

20. Lambert PP, Doriaux M, Senesael J, Vanholder R, Lammens-Verslijpe M: The pathogenicity of cationized albumin in the dog. In Lambert PP, Bergmann P, Beauwens R (Eds): *The Pathogenicity of Cationic Proteins*. New York, Raven Press, 1983, pp 307-317

21. Jeffries WA, Brandon MR, Hunt SV, Williams AF, Gatter KC, Mason DY: Transferrin receptor on endothelium of brain capillaries. *Nature* 1984; 312:162-163

22. Juranaka PF, Zastawny RL, Ling V: P-glycoprotein: Multi-drug resistance and a superfamily of membrane-associated transport proteins. *FASEB J* 1989; 3:2583-2592

23. MacDonald PN, Bok D, Ong DE: Localization of cellular retinol-binding protein and retinol-binding protein in cells comprising the blood-brain barrier of rat and human. *Proc Natl Acad Sci USA* 1990; 87:4265-4269

24. Papadrikopoulou A, Frey A, Gassen HG: Cloning and expression of γ -glutamyl transpeptidase from isolated porcine brain capillaries. *Eur J Biochem* 1989; 183:693-698

25. Solhonne B, Gros C, Pollard H, Schwartz J-C: Major localization of aminopeptidase M in rat brain microvessels. *Neuroscience* 1987; 22:225-232

26. Pardridge WM, Yang J, Eisenberg J: Blood-brain barrier protein phosphorylation and dephosphorylation. *J Neurochem* 1985; 45:1141-1147

27. Whelan JP, Eriksson U, Lampson LA: Expression of mouse β_2 -microglobulin in frozen and formaldehyde-fixed central nervous tissues: Comparison of tissue behind the blood-brain barrier and tissue in a barrier-free region. *J Immunol* 1986; 137:2561-2566

28. Barnwell JW, Asch AS, Nachman RL, Yamaya M, Aikawa M, Ingravall P: A human 88-kD membrane glycoprotein (CD36) functions in vitro as a receptor for cytoadherence ligand on *Plasmodium falciparum*-infected erythrocytes. *J Clin Invest* 1989; 84:765-772

29. Seuberger H, Lottspeich F, Risau W: The inducible blood-brain barrier specific molecule HT7 is a novel immunoglobulin-like cell surface glycoprotein. *EMBO J* 1990; 9:2151-2158

30. Krause D, Vatter B, Dermietzel R: Immunocytochemical and immunocytochemical characterization of a novel monoclonal antibody recognizing a 140 kDa protein in cerebral pericytes of the rat. *Tissue Res* 1988; 252:543-555

31. Sternberger NH, Sternberger LA: Blood-brain barrier protein recognized by monoclonal antibody. *Proc Natl Acad Sci USA* 1987; 84:8169-8173

32. Weiler-Guttler H, Sommerfeldt M, Papadrikopoulou A, et al: Synthesis of apolipoprotein A-I in pig brain microvascular endothelial cells. *J Neurochem* 1990; 54:444-450

33. Boado RJ, Pardridge WM: Molecular cloning of the bovine blood-brain barrier glucose transporter cDNA and demonstration of phylogenetic conservation of the 5'-untranslated region. *Mol Cell Neurosci* 1990; 1:224-232

34. Pardridge WM, Eisenberg J, Yang J: Human blood-brain barrier insulin receptor. *J Neurochem* 1985; 44:1771-1778

35. Weiler-Guttler H, Zinke H, Mockel B, Frey A, Gassen HG: cDNA cloning and sequence analysis of the glucose transporter from porcine blood-brain barrier. *Biol Chem Hoppe Seyler* 1989; 370:467-473

36. Pardridge WM, Boado RJ, Farrell CR: Brain-type glucose transporter (GLUT-1) is selectively localized to the blood-brain barrier. Studies with quantitative Western blotting and in situ hybridization. *J Biol Chem* 1990; 265:18035-18040

37. Pardridge WM, Eisenberg J, Yang J: Human blood-brain barrier transferrin receptor. *Metabolism* 1987; 36:892-895

38. Boado RJ, Pardridge WM: The brain-type glucose transporter mRNA is specifically expressed at the blood-brain barrier. *Biochem Biophys Res Commun* 1990; 166:174

39. Farrell CL, Pardridge WM: Blood-brain barrier glucose transporter is asymmetrically distributed on brain capillary endothelial luminal and abluminal plasma membranes—An electron microscopic immunogold study. *Proc Natl Acad Sci USA* 1991; 88:5779-5783

40. Farrell CL, Yang J, Pardridge WM: GLUT-1 glucose transporter is present within apical and basolateral membranes of brain epithelial barriers and in endothelial barriers with and without tight junctions. *J Histochem Cytochem* 1992; 40:193-199

41. Festermacher JD, Cowles AL: Theoretic limitation of intracarotid infusions in brain tumor chemotherapy. *Cancer Treat Rep* 1977; 61:519-526

42. Groothuis DR, Fischer JR, Lapin G, et al: Permeability of different experimen-

tal brain tumor models to horseradish peroxidase. *J Neuropathol Exp Neurol* 1982; 41:164-185

43. Neuwelt EA, Barnett PA, Bigner DD, et al: Effects of adrenal cortical steroids and osmotic blood-brain barrier opening on methotrexate delivery to gliomas in the rodent: The factor of the blood-brain barrier. *Proc Natl Acad Sci USA* 1982; 79:4420-4423

44. Neuwelt EA, Barnett PA, McCormick CI, et al: Osmotic blood-brain barrier modification: Monoclonal antibody, albumin, and methotrexate delivery to cerebrospinal fluid and brain. *Neurosurgery* 1985; 17:419-423

45. Neuwelt EA, Hill SA, Grenkel EP: Osmotic blood-brain barrier modification and combination chemotherapy: Concurrent tumor regression in areas of barrier opening and progression in regions distant to barrier opening. *Neurosurgery* 1984; 15:362-366

46. Tomiwa K, Hazama F, Mikawa H: Neurotoxicity of vincristine after osmotic opening of the blood-brain barrier. *Neuropathol Appl Neurobiol* 1983; 9:345-354

47. Neuwelt EA, Rapoport SI: Modification of the blood-brain barrier in the chemotherapy of malignant brain tumors. *Fed Proc* 1984; 43:214-219

48. Nagakawa H, Groothuis D, Blasberg RG: The effect of graded hypertonic intracarotid infusions on drug delivery to experimental RG-2 gliomas. *Neurology* 1984; 34:1571-1581

49. Black KL, King WA, Ikezaki K: Selective opening of the blood-tumor barrier by intracarotid infusion of leukotriene C₄. *J Neurosurg* 1990; 72:912-916

50. Aharony D, Dobson P: Discriminative effect of γ -glutamyl transpeptidase inhibitors on metabolism of leukotriene C₄ in peritoneal cells. *Life Sci* 1984; 35:2135-2142

51. DeBault LE: γ -Glutamyltranspeptidase induction mediated by glial foot process-to-endothelial contact in coculture. *Brain Res* 1981; 220:432-435

52. Soter NA, Lewis RA, Corey EJ, Austen KF: Local effects of synthetic leukotrienes (LTC₄, LTD₄, LTE₄, and LTB₄) in human skin. *J Invest Dermatol* 1983; 80:115-119

53. Black KL, Hoff JT, McGillicuddy JE, Gebarski SS: Increased leukotriene C₄ and vasogenic edema surrounding brain tumors in humans. *Ann Neurol* 1986; 19:592-595

54. Baba T, Black KL, Ikezaki K, Chen K, Becker DP: Intracarotid infusion of leukotriene C₄ selectively increases blood-brain barrier permeability after focal ischemia in rats. *J Cereb Blood Flow Metabol* 1991; 11:638-643

55. Cancilla PA, Frommes SP, Kahn LE, DeBault LE: Regeneration of cerebral microvessels: A morphologic and histochemical study after local freeze-injury. *Lab Invest* 1979; 40:74-82

56. Cancilla PA, DeBault LE: Freeze injury and repair of cerebral microvessels. In Eiserling HM, Suddith RL (Eds): *The Cerebral Microvasculature. Investigation of the Blood-Brain Barrier*. New York, Plenum, 1980, pp 257-270

57. Cancilla PA, Berliner JA, Bready JV: Astrocytes and the blood-brain barrier. Kinetics of astrocyte activation after injury and induction effects on endothelium. In Johansson BB, Owman C, Widner H (Eds): *Pathophysiology of the Blood-Brain Barrier*. Amsterdam, Elsevier, 1990, pp 31-39

58. Beck DW, Hart MN, Cancilla PA: The role of the macrophage in microvascular regeneration following brain injury. *J Neuropathol Exp Neurol* 1983; 42:601-614

59. DeBault LE, Cancilla PA: γ -Glutamyl transpeptidase in isolated brain endothelial cells: Induction by glial cells in vitro. *Science* 1980; 207:635-645

60. Maxwell K, Berliner JA, Cancilla PA: Induction of γ -glutamyl transpeptidase in cultured cerebral endothelial cells by a product released by astrocytes. *Brain Res* 1987; 410:309-314

61. Estrada C, Bready JV, Berliner JA, Pardridge WM, Cancilla PA: Astrocyte growth stimulation by a soluble factor produced by cerebral endothelial cells in vitro. *J Neuropathol Exp Neurol* 1990; 49:539-549

62. Minakawa T, Bready J, Berliner J, Fisher M, Cancilla PA: In vitro interaction of astrocytes and pericytes with capillary-like structures of brain microvessel endothelium. *Lab Invest* 1991; 65:32-40

63. Pardridge WM, Oldendorf WH, Cancilla PA, Frank HJL: Blood-brain barrier: Interface between internal medicine and the brain. *Ann Intern Med* 1986; 105:82-95

64. Beck DW, Vinters HV, Hart MN, Cancilla PA: Glial cells influence polarity of the blood-brain barrier. *J Neuropathol Exp Neurol* 1984; 43:219-224

65. Cancilla PA, DeBault LE: Neutral amino acid transport properties of cerebral endothelial cells in vitro. *J Neuropathol Exp Neurol* 1983; 42:919-929

66. Stewart PA, Wiley MJ: Developing nervous tissue induces formation of blood-brain barrier characteristics in invading endothelial cells: A study using quail-chick transplantation chimeras. *Dev Biol* 1981; 84:183-192

67. Janzer RC, Raff MC: Astrocytes induce blood-brain barrier properties in endothelial cells. *Nature* 1987; 325:253-257

68. Tao-Cheng JH, Brightman MW: Development of membrane interactions between brain endothelial cells and astrocytes in vivo. *Int J Dev Neurosci* 1988; 6:25-37